

Immune modulation of biologic systems in renal somatic cells

JERRY YEE and ERIC G. NEILSON

Renal-Electrolyte Division of the Department of Medicine, the Penn Center for Molecular Studies of Kidney Diseases and the Immunology and Cell Biology Graduate Groups, University of Pennsylvania School of Medicine, Philadelphia, Pennsylvania, USA

Components of the classical immune response mediate most traditional acts of destructive inflammation in body tissues. These events normally begin with the presentation of new peptide antigens to helper T lymphocytes by antigen-presenting cells residing within the lymphoid system. This engagement results in the elaboration of new lymphokines, and/or the emergence of new effector T cells and antibodies. These same developmental processes and events are also thought to occur when antigens are first recognized in the renal parenchyma. Not all immunologic responses within the kidney, however, are necessarily inflammatory.

The biology of somatic renal cells, for example, may only be phenotypically altered after contact with the immune system, and such contact may take place in the absence of any obvious injury. The immune response, therefore, may sometimes largely act as a signaling messenger for biologic change. Antibodies or antigen-binding soluble factors mediating these biologic changes would be specifically targeted, theoretically, to ligands expressed on certain cell types. These immune mediators are highly focused in their interaction and, depending on the second messenger signals, would produce phenotypic alterations in the target cell which are either protective or detrimental to the host. Some of the mechanisms whereby the immune system can engage and biologically alter renal somatic cells are the focus of the following review.

Antigen-binding T cell proteins

The characterization of soluble antigen-binding proteins or T-helper factors (ThF) provides a predictable mechanism for the induction of effector T cell responses in the absence of cell-to-cell contact. Such factors are mature, soluble end-products which can functionally substitute for their respective parent lymphocytes [1, 2]. They are produced by T cells, are usually antigen-specific, and traditionally engage the effector limb of an immune response [3–8]. Both T and B cell differentiation may be enhanced by such factors [9]. In contrast, antigen-binding suppressor proteins may do the opposite [9–12]. Previous biochemical characterizations of various helper factors have established that these molecules are commonly composed of two chains. One chain functions as an antigen-binding domain while the other expresses class II MHC determinants, thereby providing immunologic restriction [2, 3, 7, 9, 13–16].

At least one ThF has been described in experimental interstitial nephritis [2]. This factor induces the nephritogenic cell-mediated effector response in anti-tubular basement membrane (α TBM) disease, demonstrates fine antigenic specificity, and is

composed of two molecular chains [2, 16]. The ThF is derived from specific CD4⁺ T-helper cells reactive to 3M-1 antigen, the 30,000 M_r glycoprotein autoantigen of α TBM disease [17].

In this experimental model, genetically susceptible SJL mice (H2^s, IgH-1b) develop tubulointerstitial nephritis after immunization with renal tubular antigen in complete Freund's adjuvant [18–29]. The 3M-1 target antigen is synthesized by proximal tubular cells (MCT) and in vivo is later distributed along cortical tubular basement membranes [17, 19, 20, 21]. CD4⁺ T helper cells that are both 3M-1 antigen-specific and I-A restricted induce the appearance of CD8⁺ nephritogenic effector cells and polyclonal anti-3M-1 antibodies [18, 22, 23, 26–30].

ThF is secreted by the CD4⁺ helper cell and can be isolated as the non-covalently bound chains of a heterodimeric molecule. One chain binds 3M-1 and this antigen-binding chain expresses components of the T cell receptor (TcR) as evidenced by its ligation by several anti-T cell receptor-like antibodies [2]. Monoclonal antibody 14-30 also binds the framework region of the ThF at a determinant common to other antigen-binding factors, while another monoclonal antibody α M30 recognizes a portion of the α : β heterodimeric TcR [27]. Other antibodies which bind to this chain include KJ23, an antibody specific for V β 17, and an anti-idiotypic antibody raised against antibodies eluted from nephritis mice [31]. KJ23 recognizes a domain shared by the 3M-1 idiotype network which includes a soluble T cell suppressor factor, and effector cells [32–33]. The second chain of the ThF dimer (M_r ~82,000) expresses A β determinants, the murine equivalents of human class II MHC molecules, and binds to an anti-A β ^s monoclonal antibody.

ThF performs two known functions, the first is immunological and the second is biological. In the first, ThF promotes a dose-dependent induction of CD8⁺ effector cells, in the presence of IL-2 and 3M-1 [2]. Notably, the immunologic helper function of ThF is not perturbed even after physical disruption of its component chains through chemical alterations of alkylation and reduction [2]. The second and more novel function of this antigen-binding protein is its ability to selectively alter molecular and cellular events within proximal tubular cells.

MCT cells in culture, as an example, secrete several collagens into their supernatants, including collagen types I, III and IV [34, 35]. Culture of these cells with various concentrations of ThF leads to differential effects on the secretion of types I and IV collagen into the supernatants [36]. After 72 hours of culture, a 20-fold dose-dependent decrement in type IV collagen secretion can be observed. In contrast, the secretion of type I collagen is either unchanged or slightly enhanced by ThF [36].

The decrement in type IV collagen secretion strikingly contrasts the increases seen after the incubation of MCT cells with other cytokines and cytokine-like paracrine modulators, including TGF β , EGF and other antigen-specific T cell helper factors [2, 26].

The mechanism by which ThF down-regulates type IV collagen secretion is partially understood. In vitro experiments measuring the relative abundance of transcripts encoding collagen chains α 1(IV) and α 2(I) in MCT cells after stimulation with ThF revealed a threefold reduction in mRNA encoding the α 1(IV) chain [36]. By contrast, a slight elevation in the number of transcripts occurs with type I collagen. In vitro nuclear run-off studies demonstrate that the principal reason for the reduction in levels of mRNA encoding type IV collagen can be attributed to an overall decrease in its rate of transcription. In renal tissues of mice with early α TBM disease, T cells that stain for 3M-1 antigen-reactive ThF are found within the interstitial spaces. In situ hybridization studies of such nephritic kidneys also demonstrate a parallel reduction in transcripts encoding type IV collagen [36]. This decrease in transcripts is reflected in lower amounts of type IV collagen in basement membrane as interstitial injury progresses.

The primary constituents of tubular basement membrane are type IV collagen and laminin. The renal tubular epithelium sits on this laminar support, suspended by a fibrillar collagen scaffolding that partitions the compartments of the tubulointerstitium [37–39]. One consequence of releasing antigen-binding proteins like ThF into the interstitium is a substantial decrease in type IV collagen. Alterations in the cellular expression of extracellular matrix collagens by ThF-like moieties can directly facilitate the deformative remodeling of the tubulointerstitium during interstitial inflammation, and may provide an immunologic basis for the process of tubular atrophy and fibrogenesis [34, 36, 40, 41].

Antibody against renal cells

Antibodies to Heymann nephritis antigen complex

A paradigm for the modulating influence of antibody against antigen on glomerular epithelial cells (GEC) is Heymann nephritis, an experimental model of membranous nephropathy [42]. In the active form of disease, immunization of rats with whole kidney homogenates in adjuvant produces a heavy proteinuria. In the passive form, proteinuria is induced by the injection of either autologous or heterologous antibodies or F(Ab) $_2$ fragments that recognize a cellular antigen called Fx1A/HNAC/gp330, currently referred to as the Heymann nephritis antigen complex (HNAC) [personal communication, Dr. M.G. Farquhar; 43–47]. In both the active and passive forms of the disease, subepithelial immune deposits form along the subepithelial aspects of the GBM [46–50]. The interaction of antibody with HNAC in the tuft is not directly injurious to the GBM, since proteinuria occurs only after local complement activation; however, administration of passively transferred antibodies against brush border antigens can invoke changes along the proximal tubule in the absence of complement activation [51].

More recently, it has been observed that the antigen complex recognized by anti-HNAC antibodies (α HNAC-Ab) are likely subunits of a large, 400 to 600,000 M $_r$ multimeric membrane-associated glycoprotein (HNAC) that is found with restricted

polarity on the brush border of proximal tubular cells in the clathrin-rich inter-microvillar domain of the cell membrane and in the glomerular epithelium at the soles of the foot processes where the cell membrane faces the GBM [52–58].

Immunohistochemical localization of HNAC to the glomerular epithelial cell (GEC) has caused a reappraisal of the pathogenic mechanism of this disorder. Originally, it had been presumed that circulating immune complex formation occurred after circulating α HNAC-Ab became bound to small amounts of tubular brush border antigen that had spontaneously leaked from the kidney and entered the serum [43]. These complexes would deposit within the GBM in a subepithelial distribution and initiated disease. Currently, however, it is believed that the formation of immune deposits occurs largely in situ [37, 50, 59]. α HNAC-Ab somehow traverse the GBM and then bind avidly to HNAC in the clathrin-coated pits at the base of podocytes and subsequently form immune deposits [60–63]. The association of HNAC with clathrin, its partial similarity to brushin [58], and its homology to the LDL receptor suggests that the true nature of the complex, multimeric molecule is still uncertain [58, 64, 65]. The association of antibody with HNAC is highly specific, however, and certain epitopes are capable of inciting greater nephritogenic responses than others. For example, polyclonal brush border antibody preparations with high reactivity for HNAC produce greater proteinuria than isolated monospecific fractions [60, 66, 67]. Within these heterogeneous preparations, gp330-specific antibodies have been shown to be the most nephritogenic [58]. Cell-surface binding of α HNAC-Ab serves as the nidus for the subsequent development of subepithelial immune complexes which are shed into the lamina rarae externae and subsequently stabilized by crosslinking [47]. Aggregation of immune complexes leads to formation of large immune deposits which take form within 15 minutes and may persist in the GBM for up to eight days [47]. In culture, clustering and shedding of immune deposits occurs preferentially in the presence of polyclonal immune serum, rather than monoclonal serum, indicating that binding of multiple epitopes may occur. This arrangement likely optimizes lattice formation for the ensuing complement activation which occurs in Heymann nephritis [60, 67–69]. The enlargement of immune deposits is dependent on the continued biosynthesis of gp330 which is transported from the endoplasmic reticulum and Golgi apparatus to sites of immune interaction near the clathrin-coated pits [58]. Persistence of the immune deposits in the lamina rara externa of the GBM may occur because of ongoing exposure of nephritogenic epitopes to α HNAC-Ab within the immune complexes. Furthermore, the association of gp330 with clathrin is so strong that it is maintained even after isolation of brush border membranes by biochemical fractionation [58, 61].

With persistence of immune deposition in Heymann nephritis, significant thickening of the GBM proceeds [70]. Progressive thickening follows remodeling of the GBM structure itself [71–74]. However, the remodeling process is unusual in that there is no obvious increase in the synthesis in type IV collagen, laminin, or fibronectin [70]. Measurement of mRNA levels for these matrix components does not increase when measured from isolated glomeruli at either one week, three weeks, or three months [70]. Whether there is an increase in some other

undefined structural matrix constituent is not known. Candidates for such a component would include entactin and proteoglycans. Alternatively, a volumetric increase in the GBM may have taken place without concomitant new matrix synthesis, namely, the density of the membrane becomes decreased [70].

The nephritogenicity of HNAC may depend, in part, on expression of conformational epitopes not normally visible in vivo. After ligation by polyclonal antibodies, but not monoclonal antibodies, HNAC undergoes antigenic modulation [47, 48]. HNAC is subsequently redistributed on the apical membrane surface, patched, actively capped, and temporarily internalized after initial cross linking [47, 48]. These processes are influenced by cytoskeletal components and may be linked to one another, affecting the differential expression of epitopic specificity.

The cytoskeleton of a cell supports and provides its infrastructure. It operates dynamically and responsively to the microenvironment, in conjunction with a cell's molecular motors. As an example, the cytoskeleton behaves relatively passively when monovalent Fab' fragments of α HNAC-AB bind to Heymann nephritis antigens on the cell surface [45]. However, after binding of α HNAC-Ab and crosslinking of HNAC, the cytoskeleton becomes engaged in new actions. HNAC is redistributed and bound by α HNAC-Ab that form aggregates or "caps." These subsequently redistribute within the cell membrane [60, 75]. This capping phenomenon, which is an active process dependent on the presence of divalent cations, can be precluded by pretreatment of GEC with cytochalasin B which disrupts microfilaments, or by the addition of a calcium ionophore [75, 76]. Like capping, antigenic redistribution is also sensitive to the administration of colchicine which disrupts microtubular structures [77]. After cross linking of HNAC by α HNAC-Ab, these antigen-antibody complexes subsequently disappear from the cell membrane of the podocyte. Their subsequent reappearance in the subepithelial space at sites of slit diaphragms implies some sort of translocation to this region [47, 48].

Using differential extraction techniques, it has been shown that a conversion of HNAC from the stage of a detergent-soluble, membrane-associated form to an insoluble cytoskeleton-bound one occurs in the presence of α HNAC-Ab [76]. In the latter conformation, HNAC is tightly associated with microfilaments, thereby gaining mobility within the cytoplasm. Although the exact role of the association of the cytoskeleton with HNAC is not defined by these observations, it is likely that its role is fundamental to further antigen processing, since the administration of chlorpromazine in vitro and in vivo completely abrogates the development of Heymann nephritis [75]. Chlorpromazine inhibits the actions of calmodulin and protein kinase C and mediates a decrease in the fluid nature of the cell membrane through its effects on microfilaments [75].

The induction of proteinuria by monoclonal α HNAC-Ab is rapid, reversible and transient. This interaction is specific and does not proceed when a monoclonal antibody directed against a smaller 70,000 M_r subunit of the 600,000 M_r multimer is substituted for polyclonal α HNAC-Ab [55]. This latter antibody may interact synergistically with HNAC by increasing the permeability of the endothelium, thereby permitting a greater quantity of α HNAC-Ab to traverse the GBM, gaining access to the surface of GEC [60]. Furthermore, polyclonal α HNAC-Ab

is not uniquely nephritogenic in Heymann nephritis; a distinct antibody directed against the 108,000 M_r subunit of the 600,000 M_r multimer seems to elicit an even more severe nephritogenic response than polyclonal α HNAC-Ab [77]. Unfortunately, the molecular mechanism which governs the alteration of glomerular permeability in Heymann nephritis remains uncertain.

Interestingly, recent data also suggest a role for α HNAC-Ab in the alteration of cell adhesion of proximal tubular epithelium to the extracellular matrix. Proximal tubular epithelium in primary culture attach to their substratum and propagate as a differentiated structure [78]. These effects are mediated, in part, by HNAC. The detergent-solubilized HNAC fraction of freshly isolated epithelium has specific affinity for extracellular matrix components, including type I collagen, laminin and fibronectin, which is reminiscent of cell receptor β 1 integrins [79-81]. However, the affinity for these matrix components is neither blocked by the tripeptide sequence, RGD (arginine-glycine-asparagine), nor by synthetic fibronectin peptides, implying a non-integrin binding mechanism for HNAC. This is not unexpected since the distribution of HNAC on glomerular epithelium differs from that of the β 1 integrins. The inhibition of attachment of this epithelium to these matrix constituents in vitro by α HNAC-Ab strongly contends that a receptor epitope of HNAC mediates part of this adhesion [78]. Furthermore, the inhibition is not secondary to a toxic effect since the epithelial attachment to these matrix components is fully restored after removal of antibody from the cell surface. Another immunobiological effect of α HNAC-Ab is the prevention of the expected increase in thymidine incorporation by tubular epithelium after their culture on extracellular matrices [78].

A disturbance of proximal tubular endocytosis in rats is also mediated by α HNAC-Ab. After the injection of heterologous HNAC, there is a pronounced diminution in the endocytic uptake of FITC-conjugated dextran and horseradish peroxidase [82]. Loss of microvillar structures and flattening of the epithelial cells occurs [58]. Immunocytochemical analysis discloses a coincident and parallel reduction in the amount of several apical surface antigens, including HNAC, clathrin and an ATP-dependent proton pump. Clathrin and antigen-antibody complexes can be found in the tubular lumen. Since clathrin-coated pits play a major role in the endocytosis of filtered proteins in proximal tubules, it is not improbable that the latter phenomenon contributes to the tubular dysfunction which attends Heymann nephritis.

Antibodies to other glomerular epithelial antigens

Several other experimental models may also provide insight into potential signaling mechanisms using immunoglobulin recognition of glomerular epithelial cells. In one model in rats, a single intravenous injection of the monoclonal antibody 5-1-6 (α 5-1-6) predictably produces an immediate massive proteinuria in a dose-dependent fashion [83]. The inciting antigen has been localized exclusively to rat kidney and defines the organ specificity of this interaction. Protein excretion peaks after eight days and then gradually declines. Immunoperoxidase studies demonstrate a linear binding pattern along glomerular capillary walls within two hours after injection of α 5-1-6. By the third day, this pattern is succeeded by a fine granular pattern of fluorescence whose intensity diminishes by twelve days. This

change in immunofluorescence implies mobility of the autoantigen and quite plausibly, molecular rearrangement of immune complexes. Correlative ultrastructural studies have noted antibody *within* multivesicular bodies and localization to the surface of foot processes at slit diaphragms, similar to HNAC. The intracellular presence of antibody contrasts the mechanism of induction of Heymann nephritis where much of the antibody is seemingly not internalized. Histologically, there is partial retraction of epithelial foot processes. Taken collectively, these data imply that the determinants recognized by $\alpha 5$ -1-6 directly participate in the maintenance of the permselective function of the glomerular capillary walls. Moreover, the structural alterations therein might initiate or augment the mechanism(s) causing proteinuria. The lesions produced by $\alpha 5$ -1-6 recapitulate the histopathological findings attendant in other experimental models of non-inflammatory glomerulonephritis to include nephrosis induced by adriamycin, systemic protein overload and by the aminonucleoside puromycin [84-91].

In the K9/9 nephritis model, even more impressive structural alterations in GEC appear after the administration of this monoclonal IgG antibody. This immunoglobulin recognizes a specific sialoglycoprotein in Lewis rat kidney which leads to rapid and reversible proteinuric response. K9/9 immunoprecipitates *both* a 115 kD sialoglycoprotein of the glomerular cell membrane and a 107,000 M_r membrane component from proximal tubules [92]. Leukocytes and complement are not involved in the mechanism of proteinuria; however, there is an absolute requirement of the nonselective activation of the immune system by adjuvant for expression of the response. The structural alterations which appear include microvillous transformation, effacement of foot processes, vacuolar transformation and epithelial cell detachment from the GBM [92]. This model defines a specific nephritogenic epitope since antibodies which bind to spatially close but different epitopes cannot elicit a similar nephritogenic response [93].

In both of these models, it appears that immune recognition of epithelial ligands interferes with the orderly contact and interface of structural elements which comprise normal tissue planes. Antibody engagement subsequently disrupts the functional activity of these elements in their support of the microenvironment.

Antibodies to proximal tubular cells

Antibodies to determinants expressed by proximal tubular cells can also have an effect on MHC class II expression [94]. In α TBM disease, for example, susceptible strains of SJL mice (H-2^s, IgH-1b) develop both class I-restricted CD8⁺ and class II-restricted CD4⁺ effector cells early in disease [17, 22, 23, 26, 28-30]. Within several days, however, class I-restricted effector cells emerge as the predominant phenotype, and adoptive transfer of these particular effectors will produce disease [24].

By contrast, the non-susceptible strain of B10.A (H-2^a, IgH-1b) mice develops a class II-restricted CD4⁺ effector cell response [24]. Although this effector cell phenotype recognizes 3M-1 in delayed-type hypersensitivity assays, it is normally not nephritogenic. Such class II-restricted cells, however, can mediate tubulointerstitial inflammation upon adoptive transfer into syngeneic recipients whose tubular epithelium have been primed by an appropriate cytokine. For example, pretreatment of B10.A mice with γ IFN augments the expression of class II

antigens on proximal tubular epithelium, rendering this strain susceptible to interstitial nephritis [94]. Prior to such treatment, class II antigens are present in insufficient quantities on 3M-1-secreting tubular cells to mediate the T cell recognition of 3M-1. This is not surprising, since the normal expression of class II by proximal tubular cells is minimal [94-96]. However, under appropriate conditions, tubular epithelium can present antigen [25, 94-97], and we believe that γ IFN permissively increases the immunologic visibility of 3M-1, thus enhancing its intrinsic antigen-presenting capacity.

It appears that the permissive effect of γ IFN can be abrogated by the passive transfer of antibodies to 3M-1 [94]. B10.A mice pretreated with α 3M-1-Ab and γ IFN 48 hours before receipt of effector CD4⁺ cells prevented induction of a nephritogenic response [94]. Protection was afforded by α 3M-1-Ab even when administered as late as 24 hours before adoptive transfer. Immunofluorescent staining of class II molecules on the tubules from B10.A mice was minimal in comparison to mice which were exposed to an irrelevant control antibody, where the latter stained abundantly for class II. Additional studies suggested that the protective effect of α 3M-1-Ab was not simply due to the interference of T cell recognition of antigen, since α 3M-1-Ab could not provide a protective response in syngeneic recipients of adoptively transferred 3M-1-reactive, CD8⁺ effector cells.

Kidneys from recipients of passively transferred α 3M-1-Ab also stain TBM with increased intensity when overlain with additional α 3M-1-Ab at the time of immunofluorescent analysis. Taken together, these data suggest that passively transferred α 3M-1-Ab neither completely blocked T cell epitope recognition, nor influenced the expression of class I determinants [94].

Further studies have examined the mechanism by which α 3M-1-Ab down-regulates class II. These investigations reveal a diminution in levels of mRNA transcripts for class II determinants in MCT cells incubated with α 3M-1-Ab for 24 hours. Similar effects were obtained when polyclonal IgG eluates from kidneys of susceptible B10.S and non-susceptible B10.A strains were incubated with MCT cells. Kidney-bound α 3M-1-Ab from either strain down-regulated the transcript levels for class II antigens. Unexpectedly, however, exposure of MCT cells to the haplotype specific anti-class I antibody α K^s nearly tripled class II mRNA levels [94].

Nuclear run-off transcription assays performed on MCT cells cultured in the presence of α 3M-1-Ab pinpoint the modulatory effect of α 3M-1-Ab to an inhibition of class II gene activation [94]. A 50% decrease in the rate of class II mRNA transcription was observed with α 3M-1-Ab in comparison to levels obtained from cells incubated with control antibody. There was no effect on class I transcription. In situ hybridization studies of kidneys of mice immunized to produce α TBM disease corroborated the in vitro findings. At two or five weeks, the period during which α 3M-1-Ab are first deposited along the tubular basement membrane and before there is a strong mononuclear reaction, one-sixth to one-third of the mRNA levels of class II MHC are expressed by mice immunized to produce disease in comparison to controls [94]. Furthermore, the tubules from mice with α TBM disease demonstrate a much heavier deposition of endogenous α 3M-1-Ab than those from controls.

In summary, α 3M-1-Ab binding to proximal tubular cell surface 3M-1 selectively down-regulates the expression of class

II MHC determinants by repressing transcription. The functional and biological effects result in a diminution of cell surface class II expression that attenuate CD4⁺-mediated T cell responses. Presumably the lack of a response is the result of a failure in antigen presentation by the proximal tubular epithelium.

Summary

The various theories discussed here suggest that somatic renal cells are susceptible to biologic modulation by the immune system independent of an inflammatory effect. (1) The mode of repression of type IV collagen synthesis by novel, soluble antigen-binding proteins, the down-regulation of class II MHC expression with interruption of antigen presentation to epithelia after selective gene regulation by antibody, and the diverse interactions of antibody with renal glomerular cells producing functional disturbances in endocytosis and permselectivity; (2) modification of surface-antigen composition; (3) alteration of matrix deposition, remodeling and composition; (4) biophysical perturbation of cytoskeletal and cell membrane components; (5) and lastly, alterations in cell adhesion through cell-surface alterations, all lend testimony to the richness of the signal transduction pathways in somatic cells. Although the preceding examples represent only a small fraction of those which may take place within the glomerular and tubular microenvironments, these paradigms may nevertheless serve as new models upon which one can consider the multitude of potential communications between disparate biologic systems that connect in complex organisms.

Reprint requests to Eric G. Neilson, M.D., Renal-Electrolyte Division, 700 Clinical Research Building, University of Pennsylvania, 422 Curie Boulevard, Philadelphia, Pennsylvania 19104-6144, USA.

References

- WEBB DR, KAPP JA, PIERCE CW: The biochemistry of antigen-specific T-cell factors. *Ann Rev Immunol* 1:423-438, 1983
- HINES WH, MANN RA, KELLY CJ, NEILSON EG: Biochemical and functional characterization of an antigen-specific helper T cell factor which induces the effector T cell repertoire in autoimmune interstitial nephritis. *J Immunol* 144:75-83, 1990
- APTE RN, ESHHAR Z, LOWY I, ZINGER H, MOZES E: Characteristics of a poly(LTyr, LGlu)-poly(DLAAla)-poly(LLys)-specific helper factor derived from a T cell hybridoma. *Eur J Immunol* 11:931-936, 1981
- VANDEBRIEL RJ, DE WEGER RA, LOS G, LOVEREN HV, WIEGERS GJ, OUDE WEERNINK PA, DEN OTTER W: Two specific T cell factors that initiate immune responses in murine allograft systems. *J Immunol* 143:66-73, 1989
- DEANS JP, KROWKA JF, MOSMANN T, PILARSKI LM: Antigen-specific helper factor reacts with antibody to the T-cell receptor. *J Immunol* 14:103-107, 1987
- ZHENG H, BOYER M, FOTEDAR A, SINGH B, GREENE DR: An antigen-specific helper T cell hybridoma produces an antigen-specific suppressor inducer molecule with identical antigenic fine specificity. Implications for the antigen recognition and function of helper and suppressor inducer T cells. *J Immunol* 140:1351-1358, 1989
- ZHENG H, SAHAI BM, KILGANNON P, FOTEDAR A, GREEN DR: Specific inhibition of cell-surface T-cell receptor expression by antisense oligodeoxynucleotides and its specific effect on the production of an antigen-specific regulatory T-cell factor. *Proc Natl Acad Sci USA* 86:3758-3762, 1989
- KWONG PC, TEH HS: Characterization of a Dd-specific helper factor required for the induction of cytotoxic responses to alloantigens with the use of monoclonal antibodies specific for the helper factor of the T-cell antigen receptor. *Immunology* 61:143-150, 1987
- DEANS JP, KROWKA JF, MOSMANN T, PILARSKI LM: Antigen-specific helper factor reacts with antibody to the T-cell receptor. *J Immunogen* 14:103-107, 1987
- NEILSON EG, KELLY CJ, CLAYMAN MD, HINES WH, HAVERTY T, SUN MJ, BLANCHARD N: Murine interstitial nephritis. VII. Suppression of renal injury following treatment with a soluble suppressor factor, TsF1. *J Immunol* 139:1518-1524, 1987
- KELLY CJ, CLAYMAN MD, HINES WH, NEILSON EG: Therapeutic immune regulation in experimental interstitial nephritis with suppressor T cells and their soluble factors. *Ciba Foundation Symposium* 129:73-87, 1987
- FERGUSON TA, IVERSON GM: Isolation and characterization of an antigen-specific suppressor inducer molecule from serum of hyperimmune mice by using a monoclonal antibody. *J Immunol* 136:2896-2903, 1986
- FAIRCHILD RL, KUBO RT, MOORHEAD JW: Soluble factors in tolerance and contact sensitivity to 2,4-dinitro-fluorobenzene in mice. IX. A monoclonal T cell suppressor molecule is structurally and serologically related to the α/β T cell receptor. *J Immunol* 141:3342-3348, 1989
- GUY R, ULLRICH SJ, FOO-PHILLIPS M, HATHCOCK KS, APPELLA E, HODES RJ: Antigen-specific helper function of cell-free T cell products bearing TCR V β 8 determinants. *Science* 244:1477-1479, 1989
- YAMAUCHI K, CHAO D, MURPHY DB, GERSHON RK: Molecular composition of an antigen-specific, Ly-1 T suppressor inducer factor: One molecule binds antigen and is I-J⁻; another is I-J⁺, does not bind antigen, and imparts an Igh-variable region linked restriction. *J Exp Med* 126:423-428, 1982
- IWATA M, ADACHI M, ISHIZAKA K: Antigen-specific T cells that form IgE-potentiating factor, IgG-potentiating factor, and antigen-specific glycosylation enhancing factor on antigenic stimulation. *J Immunol* 140:2534-2542, 1988
- CLAYMAN MD, MICHAUD L, BRENTJENS J, ANDRES GA, KEFALIDES NA, NEILSON EG: Isolation of the target antigen of human anti-tubular basement membrane antibody-associated interstitial nephritis. *J Clin Invest* 77:1143-1147, 1986
- MANN R, KELLY CJ, HINES WH, CLAYMAN MD, BLANCHARD N, SUN MJ, NEILSON EG: Effector T cell differentiation in experimental interstitial nephritis. I. The development and modulation of effector lymphocyte maturation by I-J⁺ regulatory cells. *J Immunol* 136:4200-4208, 1987
- CLAYMAN MD, MARTINEZ-HERNANDEZ A, MICHAUD L, ALPER R, MANN R, KEFALIDES NA, NEILSON EG: Isolation and characterization of the nephritogenic antigen producing anti-tubular basement membrane disease. *J Exp Med* 161:290-305, 1985
- NEILSON EG, SUN MJ, KELLY CJ, HINES WH, HAVERTY TP, CLAYMAN MD, COOKE NE: Molecular characterization of a major nephritogenic domain in the autoantigen of anti-tubular basement membrane disease. *Proc Natl Acad Sci USA* 88:2006-2010, 1991
- HAVERTY TP, KELLY CJ, HINES WH, AMENTA PS, WATANABE M, HARPER RA, KEFALIDES NA, NEILSON EG: Characterization of a renal tubular epithelial cell line which secretes the autologous target antigen of autoimmune interstitial nephritis. *J Cell Biol* 107:1359-1368, 1988
- HINES WH, MANN RA, KELLY CJ, NEILSON EG: Murine interstitial nephritis: IX. Induction of the nephritogenic effector T cell repertoire with an antigen-specific T cell cytokine. *J Immunol* 144:75-83, 1990
- NEILSON EG, PHILLIPS SM: Murine interstitial nephritis. I. Analysis of disease susceptibility and its relationship to pleiomorphic gene products defining both immune response genes and a restrictive requirement for cytotoxic T cells at H-2K. *J Exp Med* 155:1075-1085, 1982
- ZAKHEIM B, McCAFFERTY E, PHILLIPS SM, CLAYMAN M, NEILSON EG: Murine interstitial nephritis. II. The adoptive transfer of disease with immune T lymphocytes produces a phenotypically complex interstitial lesion. *J Immunol* 133:234-239, 1984
- KELLY CJ, ROTH DA, MEYERS CM: Immune recognition and response to the renal interstitium. *Kidney Int* 31:518-530, 1991
- CLAYMAN MD, MICHAUD L, NEILSON EG: Murine interstitial

- nephritis. VI. Characterization of the B cell response in anti-tubular basement membrane disease. *J Immunol* 134:2375-2382, 1987
27. MEYERS CM, KELLY CJ: Effector mechanisms in organ-specific autoimmunity. *J Clin Invest* 88:408-415, 1991
 28. MANN R, ZAKHEIM B, CLAYMAN M, MCCAFFERTY E, MICHAUD E, NEILSON EG: Murine interstitial nephritis. IV. Long-term cultured L3T4⁺ T cell lines transfer delayed expression of disease as I-A restricted inducers of the effector T cell repertoire. *J Immunol* 135:286-293, 1985
 29. MANN R, NEILSON EG: The autoinduction of antigen specific Lyt 2⁺ suppressor T cells diminishes the expression of interstitial nephritis in mice with antitubular basement membrane disease. *J Immunol* 136:908-912, 1986
 30. NEILSON EG, MCCAFFERTY E, MANN R, MICHAUD L, CLAYMAN M: Murine interstitial nephritis. III The selection of phenotypic (Lyt and L3T4) and idiotype(RE-Id) T cell preferences by genes in Igh-1 and H-2K characterize the cell-mediated potential for disease expression: Susceptible mice provide a unique effector T cell repertoire in response to tubular antigen. *J Immunol* 139:2242-2249, 1985
 31. KAPPLER J, WADE T, WHITE J, KUSHNIR E, BLACKMAN M, BILL J, ROEHM N, MARRACK P: A T cell receptor V β segment that imparts reactivity to a class II major histocompatibility product. *Cell* 49:263-271, 1987
 32. NEILSON EG, ZAKHEIM B: T cell regulation, anti-idiotypic immunity, and the nephritogenic immune response. *Kidney Int* 24:289-302, 1983
 33. CLAYMAN MD, SUN MJ, MICHAUD L, BRILL-DASHOFF J, RIBLET R, NEILSON EG: Clonotypic heterogeneity in experimental nephritis. Restricted specificity of the anti-tubular basement membrane B cell repertoire is associated with a disease-modifying crossreactive idotype. *J Exp Med* 167:1296-1312, 1988
 34. KUNCIO GS, NEILSON EG, HAVERTY TP: Mechanisms of tubulointerstitial fibrosis. *Kidney Int* 39:550-556, 1991
 35. NEILSON EG, JIMENEZ SA, PHILLIPS SM: Cell mediated immunity in interstitial nephritis. III. T lymphocyte mediated fibroblast proliferation and collagen synthesis: An immune mechanism for renal fibrogenesis. *J Immunol* 125:1708-1714, 1980
 36. HAVERTY TP, KELLY CJ, HOYER JR, ALVAREZ R, NEILSON EG: Tubular antigen-binding proteins repress transcription of type IV collagen in the autoimmune target epithelium of experimental interstitial nephritis. *J Clin Invest* 89:517-523, 1992
 37. KARKEVELAS G, KEFALIDES NA, AMENTA PS, MARTINEZ-HERNANDEZ A: Comparative ultrastructural localization of collagens types III, IV, V and laminin in rat uterus and kidney. *J Ultrastruct Mol Struct Res* 100:137-155, 1988
 38. TIMPL R, DZIADEK M: Structure, development, and molecular pathology of basement membranes. *Int Rev Exp Path* 29:1-112, 1986
 39. TIMPL R, AUMAILLEY M: Biochemistry of basement membranes. *Adv Nephrol* 18:59-76, 1989
 40. HEPTINSTALL RH: Pathology of end-stage kidney disease. *Am J Med* 244:656-662, 1968
 41. CARLSON EC: Morphology of isolated renal tubular and glomerular basement membranes, in *Renal Basement Membranes in Health and Disease*, edited by PRICE RG, HUDSON BG, Orlando, Academic Press, 1987, pp. 99-114
 42. HEYMANN W, HACKEL DB, HARWOOD S, WILSON SGF, HUNTER JLP: Production of nephrotic syndrome in rats by Freund's adjuvant and rat kidney suspensions. *Proc Soc Exp Biol Med* 100:660-664, 1959
 43. EDGINGTON TS, GLASSOCK RJ, WATSON JJ, DIXON FJ: Characterization and isolation of specific renal tubular antigens. *J Immunol* 99:1199-1210, 1967
 44. GLASSOCK RJ, EDGINGTON TS, WATSON JJ, DIXON FJ: Autologous immune complex nephritis induced with renal tubular antigen. II. The pathogenetic mechanism. *J Exp Med* 127:573-588, 1968
 45. SALANT DJ, MADAJO MP, ADLER S, STILMANT MM, COUSER WG: Altered glomerular permeability induced by F(ab')₂ and Fab' antibodies to rat renal tubular epithelial antigen. *Kidney Int* 21:36-43, 1981
 46. FEENSTRA K, VAN DEN LEE R, GREGEN HA, ARENDS A, HOEDEMAEKER P: Experimental glomerulonephritis in the rat induced by antibodies directed against tubular antigens. I. The natural history: A histologic and immunohistologic study at the light and the ultrastructural level. *Lab Invest* 32:235-242, 1975
 47. KERJASCHKI D, FARQUHAR MG: Immunocytochemical localization of the heyman nephritis antigen (gp330) in glomerular epithelial cells of normal Lewis rats. *J Exp Med* 157:667-686, 1983
 48. KERJASCHKI D, MIETTINEN A, FARQUHAR MG: Initial events in the formation of immune deposits in passive Heymann nephritis: gp330-anti-gp330 immune deposits form in epithelial coated pits and rapidly become attached to the GBM. *J Exp Med* 166:109-128, 1987
 49. BARABAS AZ, LANNIGAN R: Induction of an autologous immune-complex glomerulonephritis in the rat induced by intravenous injection of heterologous anti-rat kidney tubular antibody. *Br J Exp Pathol* 55:47-55, 1974
 50. VAN DAMME JC, FLEUREN GJ, BAKKER WW, VERNIER RL, HOEDEMAEKER PJ: Experimental glomerulonephritis in the rat induced by antibodies directed against tubular antigens. V. Fixed glomerular antigens in the pathogenesis of heterologous immune complex glomerulonephritis. *Lab Invest* 38:502-510, 1978
 51. NOBLE B, ANDRES GA, BRENTJENS JR: Passively transferred anti-brush border antibodies induce injury of proximal tubules in the absence of complement. *Clin Exp Immunol* 56:281-288, 1983
 52. KERJASCHKI D, FARQUHAR MG: The pathogenic glycoprotein of Heymann nephritis is a membrane glycoprotein of the renal proximal tubule brush border. *Proc Natl Acad Sci USA* 79:5557-5561, 1982
 53. MIETTINEN A, DEKAN G, FARQUHAR MG: Monoclonal antibodies against membrane proteins of the rat glomerulus. *Am J Pathol* 137:929-944, 1990
 54. NATORI Y, HAYAKAWA I, SHIBATA S: Passive Heymann nephritis with acute and severe proteinuria induced by heterologous antibody against renal tubular brush border glycoprotein gp108. *Lab Invest* 55:63-70, 1986
 55. KERJASCHKI D, NORONHA-BLOB L, SACKTOR B, FARQUHAR MG: Microdomains of distinctive glycoprotein composition in the kidney proximal tubule brush border. *J Cell Biol* 98:1505-1513, 1984
 56. SINGH AK, SCHWARTZ MM: Nephritogenicity and immunocytochemical localization of the 70kD glycoprotein subunit (gp70) of Heymann antigen. *Clin Immunol Immunopathol* 48:61-77, 1988
 57. KERJASCHKI D, HORVAT R, BINDER S, SUSANI M, DEKAN G, OJHA PP, HILLEMANS P, ULRICH W, DONNINI U: Identification of a 400-kD protein in the brush borders of human kidney tubules that is similar to gp330, the nephritogenic antigen of rat heyman nephritis. *Am J Pathol* 129:183-191, 1987
 58. KERJASCHKI D: Molecular pathogenesis of membranous nephropathy. *Kidney Int* 41:1090-1105, 1992
 59. ANDRES G, BRENTJENS JR, CALDWELL PRB, CAMUSSI G, MATSUO S: Biology of disease. Formation of immune deposits and disease. *Lab Invest* 55:510-520, 1986
 60. BRENTJENS JR, ANDRES GA: Interaction of antibodies with renal surface antigens. *Kidney Int* 35:954-968, 1989
 61. RODMAN JS, KERJASCHKI D, MERISKO E, FARQUHAR MG: Presence of an extensive clathrin coat on the apical plasmalemma of the rat kidney proximal tubule cell. *J Cell Biol* 98:1630-1636, 1984
 62. RODMAN JS, SEIDMAN L, FARQUHAR MG: The membrane composition of coated pits, microvilli, endosomes, and lysosomes is distinctive in the rat kidney proximal tubule cell. *J Cell Biol* 102:77-87, 1986
 63. DOXSEY S, KERJASCHKI D, FARQUHAR MG: A large membrane glycoprotein (gp330) is a resident of coated pits of several absorptive epithelia. (abstract) *J Cell Biol* 97:178a, 1983
 64. RAYCHOWDHURY R, NILES JL, MCCLUSKEY RT, SMITH JA: Autoimmune target in Heymann nephritis is a glycoprotein with homology to the LDL receptor. *Science* 244:1163-1165, 1989
 65. PIETROMONACO S, KERJASCHKI D, BINDER S, ULLRICH R, FARQUHAR MG: Molecular cloning of a cDNA encoding a major pathogenic domain of the heyman nephritis antigen gp330. *Proc Natl Acad Sci USA* 87:1811-1815, 1990
 66. SPECHT GRUPP-GLANDORF RJ, DE HEER E, DEKKER-NOOREN CEM, DAHA MR, VAN ES LA: Specificity and cross-reactive idiotypes of anti-gp330 autoantibodies in active Heymann nephritis. *Immunology* 70:290-295, 1990
 67. ALLEGRI L, BRIANTI E, CHATELET F, MANARA GC, RONCO P,

- VERROUST P: Polyvalent antigen-antibody interactions are required for the formation of electron dense deposits in passive Heymann nephritis. *Am J Pathol* 125:1-6, 1986
68. COUSER WG: Mediation of glomerular injury. *J Am Soc Nephrol* 1:13-29, 1990
 69. COUSER WG, STILMANT MM, LEWIS EJ: Experimental glomerulonephritis in the guinea pig. II. Ultrastructural lesions of the basement membrane associated with proteinuria. *Lab Invest* 32:46-55, 1975
 70. FOGEL MA, BOYD CD, LEARDKAMOLKARN V, ABRAHAMSON DR, MINTO AWM, SALANT DJ: Glomerular basement membrane expansion in passive heyman nephritis: Absence of increased synthesis of type IV collagen, laminin, or fibronectin. *Am J Pathol* 138:465-475, 1991
 71. FUKATSU A, MATSUO S, KILLEN PD, MARTIN GR, ANDRES GA, BRENTJENS JR: The glomerular distribution of type IV collagen and laminin in human membranous glomerulonephritis. *Hum Pathol* 19:64-68, 1988
 72. HANSCH GM, TORBOHM I, ROTHER K: Chronic glomerulonephritis: Inflammatory mediators stimulate the collagen synthesis in glomerular epithelial cells. *Int Arch Allergy Appl Immunol* 88:139-143, 1989
 73. MATSUO S, BRENTJENS JR, ANDRES G, FOIDART JM, MARTIN GR, MARTINEZ-HERNANDEZ A: Distribution of basement membrane antigens in glomeruli of mice with autoimmune glomerulonephritis. *Am J Pathol* 122:36-49, 1986
 74. TORBOHM I, SCHOENERMARK M, WINGEN AM, BERGER B, ROTHER K, HANSCH GM: C5b-8 and C5b-9 modulate the collagen release of human glomerular epithelial cells. *Kidney Int* 37:1098-1104, 1990
 75. CAMUSSI G, NOBLE B, VAN LIEW J, BRENTJENS JR, ANDRES G: Pathogenesis of passive Heymann glomerulonephritis: Chlorpromazine inhibits antibody-mediated redistribution of cell surface antigens and prevents development of the disease. *J Immunol* 136:2127-2135, 1986
 76. CYBULSKY AV, QUIQQ RJ, BADALAMENTI J, SALANT DJ: Anti-Fx1-A induces association of heyman nephritis antigens with microfilaments of cultured glomerular visceral epithelial cells. *Am J Pathol* 129:373-384, 1987
 77. GUTMANN EJ, NILES JL, MCCLUSKEY RT, BROWN D: Colchicine-induced redistribution of an apical membrane glycoprotein (gp330) in kidney tubules. *Am J Physiol* 257:C397-C407, 1989
 78. MENDRICK DL, CHUNG DC, RENNKE HG: Heymann antigen gp330 demonstrates affinity for fibronectin, laminin, and type I collagen and mediates adherence of rat proximal tubule epithelial cell adherence to such matrices in vitro. *Exp Cell Res* 188:23-35, 1990
 79. RUOSLAHTI E: Integrins. *J Clin Invest* 87:1-5, 1991
 80. HYNES D: Integrins: A family of cell surface receptors. *Cell* 48:549-554, 1987
 81. ABRAHAMSON DR, WOODS A, COUCHMAN JR, CHATHAM WW, HECK LW: Cell surface receptors for kidney basement membranes. *Am J Kid Dis* 12:78-83, 1988
 82. GUTTMAN EJ, NILES JL, MCCLUSKEY RT: Loss of antigens associated with the endocytic pathway in proximal tubules from rats with heyman nephritis. *Am J Pathol* 138:1243-1255, 1991
 83. ORIKASA M, MATSUI K, OITE T, SHIMIZU F: Massive proteinuria induced in rats by a single intravenous injection of a monoclonal antibody. *J Immunol* 141:807-814, 1988
 84. BARNES JL, RADNIK RA, GILCHRIST EP, VENKATACHALAM MA: Size and charge selective permeability defects induced in glomerular basement membrane by a polycation. *Kidney Int* 25:11-19, 1984
 85. WASHIZAWA K, ISHII K, ITOH N, MORI T, AKABANA T, SHIGEMATSU H: Morphometric changes in the glomerular anionic sites during aminonucleoside nephrosis. *Acta Pathol* 39:558-565, 1989
 86. KANWAR YS, ROSENZWEIG LJ: Altered glomerular permeability as a result of focal detachment of the visceral epithelium. *Kidney Int* 21:565-574, 1982
 87. MESSINA A, DAVIES DJ, DILLANE PC, RYAN GB: Glomerular epithelial abnormalities associated with the onset of proteinuria in aminonucleoside nephrosis. *Am J Pathol* 126:220-229, 1987
 88. FISHMAN JA, KARNOVSKY MJ: Effects of the aminonucleoside of puromycin on glomerular epithelial cells in vitro. *Am J Pathol* 118:398-407, 1985
 89. CAULFIELD JP, FARQUHAR MG: The permeability of glomerular capillaries of aminonucleoside nephrotic rats to graded dextrans. *J Exp Med* 142:61-83, 1975
 90. WEENING JJ, VAN GULDENER C, DAHA MR, KLAR N, VAN DEN WAL A, PRINS FA: The pathophysiology of protein-overload proteinuria. *Am J Pathol* 129:64-73, 1987
 91. BERTANI G, POGGI A, POZZONI R, DELAINI F, SACCHI G, MECCA G, REMUZZI G, DONATI MB: Adriamycin-induced nephrotic syndrome in rats: sequence of pathologic events. *Lab Invest* 46:16-23, 1982
 92. MENDRICK DL, RENNKE HG: I. Induction of proteinuria in the rat by a monoclonal antibody. *Kidney Int* 33:818-830, 1988
 93. MENDRICK DL, RENNKE HG: II. Epitope specific induction of proteinuria by monoclonal antibodies. *Kidney Int* 33:831-842, 1988
 94. HAVERTY TP, WATANABE M, NEILSON EG, KELLY CJ: Protective modulation of class II MHC gene expression in tubular epithelium by target antigen-specific antibodies. Cell-surface directed down-regulation of transcription can influence the susceptibility to murine tubulointerstitial nephritis. *J Immunol* 143:1133-1141, 1989
 95. JEVIKAR AM, WUTHRICH RP, TAKEI F, XU HW, BRENNAN DC, GLIMCHER LH, KELLY VE: Differing regulation and function of ICAM-1 and class II antigens on renal tubular cells. *Kidney Int* 38:417-425, 1990
 96. HINES WH, HAVERTY TP, ELIAS JA, NEILSON EG, KELLY CJ: T cell recognition of epithelial self. *Autoimmunity* 5:37-47, 1989
 97. WUTHRICH RP, GLIMCHER LH, YUI MA, JEVIKAR AM, DUMAS SE, KELLY VE: Generation of highly differentiated murine renal tubular epithelial cell lines: MHC class II regulation, antigen presentation, and tumor necrosis factor production. *Kidney Int* 37:783-792, 1990